

Go to the class website and download this file: [SDSS\\_DR7.dat](#)

It contains data from the final (Data Release 7) SDSS galaxy redshift survey. Specifically, the file contains 550,166 lines, where each line has the following info for a single galaxy:

$RA$  – right ascension  
 $DEC$  – declination  
 $z$  – redshift  
 $M_g$  – absolute  $g$ -band magnitude  
 $M_r$  – absolute  $r$ -band magnitude

**a)** Familiarize yourself with the data: Make a plot of  $DEC$  (y-axis) vs.  $RA$  (x-axis) showing the positions of the galaxies on the sky. Make a plot of  $M_r$  (y-axis) vs.  $z$  (x-axis) also showing all the galaxies (invert the y-axis so that more luminous galaxies are up).

**b)** Make a plot of the  $(g-r)$  color distribution of galaxies. Using a cut between red and blue galaxies of  $(g-r)=0.75$ , calculate the fraction of blue galaxies in the sample.

**c)** Plot the  $r$ -band luminosity function of galaxies: a histogram of  $\log(dn/dM_r)$  as a function of  $M_r$ , where  $dn$  is the number density of galaxies (in units of  $h^{-1}\text{Mpc}$ ) in bins of  $M_r$  (use bins of width 0.1). To compute the density, you need to estimate a volume. This is difficult for a flux-limited sample since it has no clear redshift boundary, so just use the median redshift survey depth of  $z<0.1$ . You will also need the area on the sky covered by the SDSS DR7, which is  $7,675.2 \text{ deg}^2$  (2.295 steradians).

**d)** Construct three volume-limited sub-samples of the full dataset that contain galaxies more luminous than -20, -19, and -18, respectively. List the redshift bounds, the volume, and the number of galaxies for each sample. Calculate new blue galaxy fractions for these samples.

**e)** Make a single plot of the luminosity functions measured from these three volume-limited samples. How do these look different from the luminosity function made using the flux-limited sample?