


# Dust Maps

# Interstellar Dust


---

- Robert Trumpler first confirmed the existence of interstellar dust (1930; see syllabus link and, in particular, point 5 of the Summary on page 187)
- Trumpler noticed that open star clusters of the same general apparent size had different brightnesses
- Either the real size of clusters increases farther from the Earth (i.e. the Earth is in a special location), or stars become progressively *fainter* at larger distances
- Absorption by interstellar dust would make stars progressively fainter with distance from the Earth



# Dust Maps

- Now, the third most cited journal paper in astronomy deals with the careful mapping of interstellar dust
- Schlegel et al. (1998; see syllabus links)
- The paper is well-cited because any measurement of the flux (or magnitude) from extragalactic sources must correct for foreground absorption by dust in our Galaxy
- To correct a magnitude for dust, subtract the extinction ( $A$ ) in that band ( $m_{true} = m_{observed} - A$ ; additional equations for interstellar reddening are linked from the syllabus)



# Dust Maps

---

- You will need to install *sfdmaps*. You can do this with *pip install sfdmap* in the command line.
  - And download the SFD maps. In your week4 directory:  
*git clone https://github.com/kbarbary/sfddata*
  - Make sure that you do not add the *sfddata/* directory to our repository tracking!
  - Now you can use the *sfdmap* package and *astropy.coordinates* to access the maps:
  - Convert (RA, Dec) to Galactic coordinates:
    - *ra, dec = '00h42m30s', '+41d12m00s'*
    - *c = SkyCoord(ra, dec).galactic*
-

# Dust Maps

---

- Obtain the reddening at this position from the dust maps
    - *import sfddmap*
    - *dustdir = './sfddata/'*
    - *m = sfddmap.SFDDMap(dustdir, scaling=1)*
    - *ebv = m.ebv(c.l.value, c.b.value, frame='galactic')*
  - Note that it is also possible to obtain the reddening without first converting to Galactic coordinates, e.g.
    - *ra, dec = '00h42m30s', '+41d12m00s'*
    - *c = SkyCoord(ra, dec)*
    - *ebv = m.ebv(c.ra.value, c.dec.value)*
-

# Dust Maps

---

- Finding *extinction* requires wavelengths and a dust-law from the *extinction* package.
  - Install the extinction package: *pip install extinction*.
  - To find *rough* extinctions for the SDSS *ugriz* filters (more on *ugriz* later):
    - *import extinction*
    - *wave = np.array([3543., 4770., 6231, 7625., 9134.])*
    - *A = extinction.fitzpatrick99(wave, 3.1\*ebv)*
-

# Python tasks

---

1. The objects at  $(\alpha, \delta) = (246.933^\circ, 40.795^\circ)$  and  $(236.562^\circ, 2.440^\circ)$  are both quasars near a redshift of  $z = 1.08$ 
    - Use the *SDSS Navigator Tool* linked from the syllabus to obtain the magnitudes of these quasars and plot  $g - r$  versus  $r - i$  for both quasars in the same plot
    - Do the quasars have similar colors? Should they?
    - Correct the quasars' magnitudes for Galactic extinction and re-plot them. Do their colors now agree better?
  2. Let's visualize the dust in the region of each quasar
    - Use *numpy.meshgrid* to make a 2-dimensional 100 x 100 array (i.e. a grid) centered near  $(236.6^\circ, 2.4^\circ)$  with  $1^\circ$  bins. Create another set centered at  $(246.9^\circ, 40.8^\circ)$  for RA in  $1.3^\circ$  bins and for DEC in  $1^\circ$  bins.
-

# Python tasks

---

3. Convert your RA/DEC 100 x 100 arrays to Galactic ( $l$ ,  $b$ )
  4. Using matplotlib's *contour* procedure (linked from the syllabus), let's plot dust maps
    - Find the amount of reddening  $E(B-V)$  at each *position* in your 100 x 100 arrays.
    - Make a contour plot:
      - $cs = plt.contour(RA, Dec, ebmV)$
      - Think about appropriate contour levels.
  5. On your plots, also show the position of each quasar and a line in RA/Dec depicting the Galactic Plane (a line from  $l = 0^\circ$  to  $360^\circ$ ;  $np.arange(360)$ ; at  $b = 0^\circ$ ;  $np.zeros(360)$ ]
-