


Time Domain Surveys: Variability

Time Domain Surveys: Variability


- Time domain science will be hot in the next decade, particularly in the era of the *Vera Rubin Observatory (VRO)*



- Movies of the sky record transient phenomena, such as explosions, bursts, quasars, transits, variable stars and flares (see the *LSST Science Book Chapter* syllabus link)
 - *VRO* will scan the sky once every few nights
 - the typical time between observations of an object in a time domain survey is called the *cadence*
-


Time Domain Surveys: Variability

- Current surveys with a time domain component include the *ZTF* and the *Dark Energy Survey* supernova effort (both linked from the syllabus)
- No current survey has a cadence and duration that is close to matching the *LSST*, but *SDSS Stripe 82* (pictured) is a useful testbed for time domain science
- The cadence of *SDSS Stripe 82* is a little strange, being a combination of annual and daily imaging runs (as we shall see) but it's illustrative of time domain information



Temporal Queries of SDSS Stripe 82

- The full *SDSS Stripe 82* database is huge (about 100 times larger than an SDSS run of comparable area)
 - so we don't have a local copy
- But, HTM indexing is rapid for this sort of application (matching multiple observations that are very close in position...i.e. multiple observations of the same object)
- So, the online *SDSS Catalog Archive Server* coupled with careful construction of an *SQL* query works well



Temporal Queries of SDSS Stripe 82

- The *SDSS Stripe 82 Catalog Archive Server* is linked from the syllabus (it's different to the one that we used in earlier lectures to query the single-epoch SDSS data)
 - To return all objects within 0.3 arcseconds of a sky position (ra , dec), at any time of observation, use, e.g.:
 - *SELECT p.ra, p.dec, p.psfmag_g, f.mjd_g, n.distance*60
FROM fGetNearbyObjEq(ra,dec,0.3/60) n, PhotoPrimary p,
Field f WHERE n.objID = p.objID and f.fieldID = p.fieldID*
 - Here, the “ n ” is the database of HTM indexes, the “ p ” is the database of imaging information for primary objects and the “ f ” database records the time that each image was observed
 - The $_g$ variables convey the fact that we only want to return magnitudes and times of observation (MJD) in the g-band
-

Temporal Queries of SDSS Stripe 82

- Code in my week12 directory can *remotely* run the query

```
SELECT p.ra, p.dec, p.psfmag_g, f.mjd_g, n.distance*60  
FROM fGetNearbyObjEq(ra,dec,0.3/60) n, PhotoPrimary p,  
Field f WHERE n.objID = p.objID and f.fieldID = p.fieldID
```
 - The code is called stripe82query.py and takes RA and Dec as arguments passed at the command line
 - The output will be the same whether you use my Python code or the online query, but my code can be spawned from other Python code using, e.g., *os.system*
 - The output can be directed to a file and read back in:
 - *os.system("python stripe82query.py 29.2256832 0.4208970 > sqlresult.data")*
-

Python tasks

1. Query the SDSS Stripe 82 database and retrieve the time domain data for $(\alpha, \delta) = (29.2256832, 0.4208970)$, $(35.3756676, 0.0017000)$, $(45.299833, -0.55386111)$, $(58.175468, 0.218697)$ and $(60.829041, -1.240793)$
 - Plot g (y-axis) against MJD (x-axis) for *all the objects on one plot*...why does the cadence looks as it does?
 - Each object may have an observation at $mjd_g = 0$ for the *run* that corresponds to the coadded data ($run = 106$ or $run = 206$)...you can remove this observation
 2. Plot MJD against g for *each individual object*
 - Try to determine which of these objects is a *supernova*, which are *normal stars*, which is a *quasar* and which is a *short-period (a few days) variable star*
-

Python tasks

3. I drew some of these objects from the *SDSS stripe 82 Variable Source* and *Standard Star* Catalogs (linked from the syllabus) so they are not necessarily representative of typical sources in *Stripe 82*

- Find a few objects in the coadded imaging file at `/astro/astr8020/varcats/varcat-ra30-60-minflux_ngood1-mincoadd_flux_ngood2.fits` that have more than 25 observations (i.e. $FLUX_NGOOD > 25$ in every band)
 - Download these objects' time domain information from the *Stripe 82 SQL Server* and plot *MJD* against *g* for each object
 - Does a typical object in *Stripe 82* have much variability (i.e. does it look like the *normal stars* from task 2)?
-